

Name : _____

Date : ____ / ____ / ____

1) How would you describe the overall level of AS neck, back or hip pain you have had?

| | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|----|
| 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|---|---|---|---|---|---|---|---|---|---|----|

None

Very severe

2) How long does your morning stiffness last from the time you wake up?

| | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|----|
| 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|---|---|---|---|---|---|---|---|---|---|----|

0

1

2 or more hours

3) How active was your spondylitis on average during the last week?

| | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|----|
| 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|---|---|---|---|---|---|---|---|---|---|----|

Not active

Very active

4) How would you describe the overall level of pain/swelling in joints other than neck, back or hips you have had?

| | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|----|
| 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|---|---|---|---|---|---|---|---|---|---|----|

None

Very severe


5) C-reactive protein (mg/L)?

| Back Pain | |
|-----------|-----|
| 0 | 0 |
| 1 | 0,1 |
| 2 | 0,2 |
| 3 | 0,4 |
| 4 | 0,5 |
| 5 | 0,6 |
| 6 | 0,7 |
| 7 | 0,8 |
| 8 | 1,0 |
| 9 | 1,1 |
| 10 | 1,2 |

| Morning Stiffness | |
|-------------------|-----|
| 0 | 0 |
| 1 | 0,1 |
| 2 | 0,1 |
| 3 | 0,2 |
| 4 | 0,2 |
| 5 | 0,3 |
| 6 | 0,4 |
| 7 | 0,4 |
| 8 | 0,5 |
| 9 | 0,5 |
| 10 | 0,6 |

| Patient Global | |
|----------------|-----|
| 0 | 0 |
| 1 | 0,1 |
| 2 | 0,2 |
| 3 | 0,3 |
| 4 | 0,4 |
| 5 | 0,6 |
| 6 | 0,7 |
| 7 | 0,8 |
| 8 | 0,9 |
| 9 | 1,0 |
| 10 | 1,1 |

| Peripheral Pain/Swelling | |
|--------------------------|-----|
| 0 | 0 |
| 1 | 0,1 |
| 2 | 0,1 |
| 3 | 0,2 |
| 4 | 0,3 |
| 5 | 0,4 |
| 6 | 0,4 |
| 7 | 0,5 |
| 8 | 0,6 |
| 9 | 0,6 |
| 10 | 0,7 |



| C-Reactive Protein (mg/L) | | | | | | | |
|--|-----|----|-----|-----|-----|-----|-----|
| Note: when the conventional CRP is below the limit of detection or when the high sensitivity CRP is <2mg/L, the constant value of 2mg/L should be used to calculate ASDAS-CRP. | | | | | | | |
| ≤2 | 0,6 | 15 | 1,6 | 60 | 2,4 | 125 | 2,8 |
| 3 | 0,8 | 16 | 1,6 | 65 | 2,4 | 130 | 2,8 |
| 4 | 0,9 | 17 | 1,7 | 70 | 2,5 | 135 | 2,8 |
| 5 | 1,0 | 18 | 1,7 | 75 | 2,5 | 140 | 2,9 |
| 6 | 1,1 | 19 | 1,7 | 80 | 2,5 | 145 | 2,9 |
| 7 | 1,2 | 20 | 1,8 | 85 | 2,6 | 150 | 2,9 |
| 8 | 1,3 | 25 | 1,9 | 90 | 2,6 | 155 | 2,9 |
| 9 | 1,3 | 30 | 2,0 | 95 | 2,6 | 160 | 2,9 |
| 10 | 1,4 | 35 | 2,1 | 100 | 2,7 | 165 | 3,0 |
| 11 | 1,4 | 40 | 2,2 | 105 | 2,7 | 170 | 3,0 |
| 12 | 1,5 | 45 | 2,2 | 110 | 2,7 | 175 | 3,0 |
| 13 | 1,5 | 50 | 2,3 | 115 | 2,8 | 180 | 3,0 |
| 14 | 1,6 | 55 | 2,3 | 120 | 2,8 | 185 | 3,0 |

